भारतीय मानक Indian Standard

मणिक्कड़ई नूल (कलाई परिधि चिन्ह) विश्लेषण — रीति संहिता

IS 19055: 2024

Maņikkaţai nūl (Wrist Circummetric Sign) Analysis — Code of Practice

ICS 11.120.10

© BIS 2024



भारतीय मानक ब्यूरो BUREAU OF INDIAN STANDARDS मानक भवन, 9 बहादुर शाह ज़फर मार्ग, नई दिल्ली - 110002 MANAK BHAVAN, 9 BAHADUR SHAH ZAFAR MARG NEW DELHI - 110002

www.bis.gov.in www.standardsbis.in

FOREWORD

This Indian Standard was adopted by Bureau of Indian Standards, after the draft finalized by Siddha Sectional Committee had been approved by the Ayush Division Council.

There is a worldwide resurgence of interest in holistic systems of health care, particularly with respect to the prevention and management of lifestyle-related disorders, and chronic, non- communicable, and systemic diseases. It is increasingly understood that no single healthcare system can provide satisfactory answers to all the health needs of modern society. Evidently, there is a need for a new inclusive and integrated healthcare regime that should guide health policies and programs in the future.

Siddha medicine is an ancient medical system in India and involves a system of regional therapy with geographical significance for the convenience of treatment and comfort of patients. The Siddha system involves the use of herbs, minerals, metals, and any organic material available, and emphasizes patients' physiology, environment, age, gender, race, habits, psychology, habitat, diet, appetite, physical condition, and disease.

In Siddha Medicine, Akattiyar cūtāmaṇi kayaru cūttiram provides diagnostic and prognostic predictions correlating with anthropometric values of wrist circumference measured by one's own finger breadths. This gives 96 descriptions under 26 finger breadth measurements and subdivisions. The Maṇikkaṭai nūl (Wrist Circummetric Sign) provides insight into Siddha Anthropometry giving analogues predicting one's current health status and prognosis as well. Standardization of this ancient techniques helps for wider use of this technique will influence ascertaining clinical outcome and physician decision.

This standard will provide the requirements for the techniques and the practice guidelines. It will help in providing guidance to Siddha Physicians, Researchers, Academicians, etc, to analyze *Maṇikkaṭai nūl* appropriately and help establish it as a traditional diagnostic tool for Preventive, Personalized and Prophylactic treatment through Siddha Medicine.

The composition of the Committee responsible for the formulation of this standard is given in Annex C.

For the purpose of deciding whether a particular requirement of this standard is complied with, the final value, observed or calculated, expressing the result of a test or analysis, shall be rounded off in accordance with IS 2:2022 'Rules for rounding off numerical values (*second revision*). The number of significant places retained in the rounded off value should be the same as that of the specified value in this standard.

Indian Standard

Maṇikkaṭai nūl (WRIST CIRCUMMETRIC SIGN) ANALYSIS — CODE OF PRACTICE

1 SCOPE

This standard specifies general requirements and code of practice details for *Maṇikkaṭai nūl* (Wrist Circummetric Sign) Analysis.

2 REFERENCES

The standards given below contain provisions which, through reference in this text, constitute provision of this standard. At the time of publication, the editions indicated were valid. All standards are subject to revision, and parties to agreements based on this standard are encouraged to investigate the possibility of applying the most recent edition of these standards:

IS No.	Title
IS 2819 : 2024	Textiles — Braided cotton cord — Specification (third revision)
IS 4513 :1968	Specification for scissors, surgical dressing and stitch
IS 1481 : 2024	Metric steel scales for engineers — Specification (second revision)

3 TERMINOLOGY

For the Purpose of this standard, the following terms and definitions shall apply.

- **3.1** *Maṇikkaṭai nūl* A unique diagnostic strategy/tool used in Siddha System of Medicine to predict certain clinical features occurring in an individual based on wrist circumference measured using an inelastic thread.
- **3.2** *Maṇikkaṭai* The term "Maṇi" denotes wrist and "kaṭai" means end measures. The measure of the wrist (circumference) obtained at the antebrachial point by using a $n\bar{u}l$.
- **3.3** $N\bar{u}l$ Suitable non elastic thread/rope with specifications as mentioned in 4.1.
- **3.4** $Viral\ katai$ The term "Viral" denotes finger and "katai" means end measures. The measure of the mediolateral breadth/width of individual finger unit using a $n\bar{u}l$.

- **3.5 Finger Breadth (FB)** One full breadth of the finger is taken into consideration for FB and the readings of each FB (unit) are divided into 4 units, quarter measurement ($\frac{1}{4}$), half ($\frac{1}{2}$), three quarters ($\frac{3}{4}$), and full breadth.
- **3.6 Calculation of Finger Breadth (FB)** Unit of measure calculating the mediolateral width of one extended finger (palm-side).
- **3.7** *Nānku viral kaṭai* (**4 Finger Breadth**) The mediolateral width of the 4-extended fingers (palm-side) of the patient.
- **3.8 Ideal point of** $N\bar{a}\underline{n}ku$ $vi\underline{r}al$ $ka\underline{r}ai$ (4 Finger Breadth) measurement The middle location of the four fingers where the $n\bar{u}l$ (thread) is placed to measure the $N\bar{a}\underline{n}ku$ $vi\underline{r}al$ $ka\underline{r}ai$.
- **3.9 Antebrachial point (ABP)** The point at the forearm/antebrachium where the circumference is measured using the $n\bar{u}l$ (thread). It is an important point in the forearm (Antebrachial region) which is used to measure the Antebrachial circumference. The length of the four finger breadths is measured back from the wrist line to reach this focus.
- **3.10 Antebrachial Circumference** (**ABC**) The circumference of ABP as measured using the $n\bar{u}l$. Further, the length obtained in the $n\bar{u}l$ is converted into number of Finger Breadth Unit (FBU) to assess readings.
- 3.11 Anatomical Point for $N\bar{a}\underline{n}ku$ viral kaṭai (4 Finger Breadth) Measurement $N\bar{a}\underline{n}ku$ viral kaṭai (4 finger breadth) as referred to in 3.8.
- **3.12** Anatomical Point for *Maṇikkaṭai* (Wrist Circummetric) Measurement Antebrachial point as referred to in **3.9**.

4 REQUIREMENTS

4.1 Cotton Yarn (Thread) Cord

The $n\bar{u}l$ (thread) used for measurement shall be made of cotton yarn uniformly braided with 3 strands. The ideal range of length of the cord shall be ranging from 150 mm to 200 mm, and width of nominal diameter around 3 mm. The $n\bar{u}l$ shall be free from knots, kinks, broken or loose ends projecting from the cord.

5 METHODS FOR MAŅIKKAŢAI NŪL (WRIST CIRCUMMETRIC SIGN)

5.1 Method 1

- a. The patient is asked to sit comfortably in the chair provided, by keeping the hand fingers intact and straight. (See Fig. 1);
- b) The patient is asked to hold the wrist anterior surface up and place it on the examination table for easy measuring by the investigators. (See Fig. 2);
- c) The specified $n\bar{u}l$ (thread) is used to take the measurement. (See Fig.3);
- d) The total mediolateral width of 4 fingers is measured from the Ideal point of *Nanku*

- *Nānku viral kaṭai* measurement by using the *nūl* (*See* Fig. 4 and Fig. 5);
- e) The length obtained, that is, ABC is used to ascertain the anti-brachial circumference spot (*See* Fig. 6, and Fig. 7), then converted into total number of FBU and recorded. (*See* Fig. 8);
- f) Repeat the process twice again for maximum accuracy of the reading. (*See* Fig. 9); and
- g) The reading obtained in number of FBU is interpreted with clinical description from the script (see Annex A).



FIG. 1 FINGERS
BROUGHT TOGETHER
AND PLACED STRAIGHT



FIG. 2 PALM PLACED FACING UPWARDS



Fig. 3 *Nūl* (Thread) for Measurement



FIG. 4 MEASURING TOTAL MEDIOLATERAL WIDTH OF 4 FINGERS



Fig. 5 Ideal Point of *Nāṇku viṛal kaṭai*



FIG 6 IDENTIFY THE ANTEBRACHIAL CIRCUMFERENCE SPOT



FIG. 7 MEASURING ANTEBRACHIAL CIRCUMFERENCE



Fig. 8 4 Finger Breadth Units Recorded



FIG. 9 4 FINGER BREADTH UNITS CHECKED WITH ONE FINGER BREADTH FOR ACCURACY

5.2 Method 2

- a) The patient is asked to sit comfortably in the chair provided, by keeping the hand fingers intact and straight. (See Fig. 1A);
- b) The patient is asked to hold the wrist anterior surface up and place it on the examination table for easy measuring by the investigators. Ask the patient to keep their fingers together (Adduction of fingers). (See Fig. 2A);
- c) The patient is asked to place left or right hand over the anterior wrist region so that the little finger is placed in the same plane of wrist line (See Fig. 3A);
- d) A point is marked on forearm where the outer border of index finger lies and that is the antebrachial circumference point. (See Fig. 4A);
- e) The antebrachial circumference is measured (ABC) by using the same $n\bar{u}l$. (See Fig. 5A);
- f) The length obtained, that is, ABC is

- converted into the total number of FBU and recorded. (*See* Fig. 6A);
- g) The process is repeated twice again for maximum accuracy of the reading; and
- h) The reading obtained in the number of FBU is interpreted with clinical description from the script. (*See* Annex A).

5.3 Calculation of *Manikkaṭai nūl* Value:

- a) The nearest measure of FBU (full breadth, three quarters, half and one-fourth) is confirmed based on the consequent reading;
- b) The findings are expressed as FBU is calculated using the formula (width of the 4 extended fingers (Palmar) of the Patient/4); and
- c) The FBU values provided in <u>Annex A</u> is based on the Classical textual references and are given solely for Physician's interpretation.




FIG. 10 ANATOMICAL POINT FOR MANIKKAŢAI MEASUREMENT




FIG. 11 ANATOMICAL PLANES OF THE FOREARM

5.4 Conditions for Measurement:

- a) While the measurement can be made on either left or right sides of the forearm, the report shall clearly indicate on which side the measurement has been taken. It is to be considered, while in *Nāṭi paricōtaṇai* (Pulse examination), generally right side is used for men and left side is used for women;
- b) Presence of any anatomical deformities, previous fractures of the arm is to be examined and noted as it may affect the accuracy of readings. If observed reading is to be taken in the opposite arm; ; and
- c) History of foreign bodies (metal rods, plates etc.) in-situ, arteriovenous (AV) fistula is to be considered.



FIG. 1 (A) FINGERS BROUGHT TOGETHER AND PLACED STRAIGHT



FIG. 2 (A) PALM PLACED FACING UPWARDS



FIG. 3 (A) PLACING RIGHT-HAND 4 FINGERS OVER THE LEFT-HAND ANTERIOR WRIST



Fig. 4 (A) Ideal Antebrachial Circumference Point



FIG. 5 (A)
MEASURING
ANTEBRACHIAL
CIRCUMFERENCE



FIG. 6 (A)
CONVERTING THE
ANTEBRACHIAL
CIRCUMFERENCE INTO
TOTAL NUMBER OF
FINGER BREADTH
UNITS

IS 19055 : 2024

6 SPECIAL CONSIDERATION

Conditions such as Congenital abnormality of the wrist, any abnormal swelling, injury or fracture of the wrist, dropsy, malnutrition, foreign bodies insitu are to be taken into consideration before establishing *Manikkaṭai* values.

7 EXPRESSIONS OF RESULT

- a) The findings are to be correlated with the presenting complaints and note of patients;
- b) Leading questions are to be asked and logical derivations used to interpret the diagnosis; and
- c) The results are analyzed based on the affected *Mukkurram* and the impacted *capta tātu*, expressed as the subjective signs, and findings as per suitable format (Annex B).

8 REPEATABILITY

The accuracy of the repeatability is obtained by at least three consecutive readings.

9 DATA COLLECTION PROFORMA REQUIREMENTS

9.1 Basic Demographic Description of Subjects

Fresh data of the patient should be collected through either case record forms (CRF) or questionnaire/case reports. CRF could be generated according to the expected specific outcome of the studies or in general. Information such as demographic details, clinical history relatable to *Maṇikkaṭai nūl* measurements and vital data on current clinical status are to be presented in the data collection proforma. A model CRF for Diagnostic cum validation studies of *Maṇikkaṭai nūl* is given in Annex B for reference.

10 OTHER CONSIDERATIONS

The Maṇikkaṭai nūl analysis can be used to analyze the state of the Mukkuṛram and capta tātu. The analysis can be more productive when the body constitution (Uṭal tātu/ Yākkai ilakkaṇam) is also taken into consideration along with other details. This analysis can be used as an effective diagnostic tool to improve the accuracy of the disease prediction.

ANNEX A

(Clauses $\underline{5.1}$, $\underline{5.2}$ and $\underline{5.3}$)

MANIKKAŢAI NŪL (WRIST CIRCUMMETRIC SIGN — INFERENCE)

SI No.	Maṇikkaṭai nūl Measurements	Inferences given in the Siddha Classical Books
	(in FB*)	
(1)	(2)	(3)
i)	11 FB	Considered a healthy sign; as otherwise, the patient may be obese;
ii)	10 FB	Diseases that are causing pallor (Anaemia); Body and extremities appear shrunken; Cluster of gastrointestinal diseases attended with pain and painful gastrointestinal disorders accompanied by indigestion.
iii)	9 ^{3/4} FB	Sinus ulcer; Inguinal lymphadenopathy; Cough; Dryness; Diseases of the spleen may occur within a year
iv)	9 ^{1/2} FB	Heat affecting the bones with the occurrence of pallor of the body; Dryness of eyes; Fever; Dryness of the body
v)	9 ^{1/4} FB	Oliguria or reduced urinary output; Insomnia due to intense sensation of heat; Sinusitis and its associated symptoms
vi)	9 FB	Pricking pain inside the ear with ear blockage or dullness;
		Pain in the hip region and thigh;
vii)	8 ^{3/4} FB	Dryness of the body; Skin diseases resembling urticaria; Pricking pain in the abdomen; <i>Pitta</i> diseases, - causing pallor of conjunctiva, oral cavity, and Upper extremity.
viii)	8 ^{1/2} FB	Cough; Sensation of heat in the body; Formation of ulcer; Glandular swelling
ix)	8 ^{1/4} FB	Febrile illness of <i>Pitta vāyu</i> nature characterized by yellowish discoloration of body, vomiting of in digested matter with colicky painof the abdomen, and dropsy.; Venereal disease (Sexual transmitted disease); Diseases of the head may occur within one year
x)	8 FB	Flatulence causing stomach growling; Swelling in the body; Chronic Sinusitis; Throat pain
xi)	7 ^{3/4} FB	Ano-rectal diseases (haemorrhoids); Weakening of both lower extremities; Tremor of the head; Enlarged lymph nodes in the cervical region (Cervical lymphadenopathy) may occur within one year, along with bleeding per nose; Over a period of time there will be discharge ofblood from the nose
xii)	7 ^{1/2} FB	Group of diseases in which there is dryness, softening, soft necrosis, caries, tuberculosis, decay or destructive changes in bones, cartilages orother bony tissues accompanied by general emaciation, muscular dystrophy and fatigue.; Flatulence and abdominal distension; Burning sensation of face; Burning sensation of eyes; Febrile illness; Pain in thelegs; Occurrence of Pyemic abscess or Carbuncle or Carcinoma in the thigh region
xiii)	7 ^{1/4} FB	Low back ache; Bilious aggression in the head region and its associated symptoms; Diseases that are causing pallor (Anaemia); Heaviness of both extremities; Insomnia
xiv)	7 FB	Bilious aggression in the head region which may present the features of per oral bleeding, Pulmonary tuberculosis and dryness of the stool associated with increased body heat

SI No.	Maṇikkaṭai nūl Measurements (in FB*)	Inferences given in the Siddha Classical Books
(1)	(2)	(3)
xv)	6 ^{3/4} FB	Swelling or inflammation of the scrotum due to the vitiation of <i>Vātam</i> ; Reddish discoloration of eyes; State of stupor or giddiness; Urinary obstruction may occur within a period of 3 years; Pain in both upper and lower extremities
xvi)	6 ^{1/2} FB	Febrile conditions or increased body heat; Pricking pain in the body; Itmay be a death sign
xvii)	6 ^{1/4} FB	Anorectal diseases like haemorrhoids; Bloody dysentery; Severe diarrhoea associated with sleeplessness; Change in the skin colour or its darkening
xviii)	6 FB	Severe phlegmatic congestion in the chest; Stupor or giddiness; It may be a fatal sign
xix)	5 ^{3/4} FB	Delirium and its associated features; Giddiness; Fatigue; It may be a fatal sign
xx)	5 ^{1/2} FB	Toxin affecting head; Pallor of the nose; Tooth becomes blackish in nature; It may be a fatal sign (Death may be predicted within 10 days)
xxi)	5 ^{1/4} FB	Severe emaciation (cachexia) of the body; Increased sleepiness; It may be a fatal sign (Death may be predicted within 6 days)
xxii)	5 FB	Pallor of the body and chillness; Signs of phlegm congestion resembles that of toxicity; It may be a fatal sign.
xxiii)	4 ^{3/4} FB	A state of unconsciousness; Tongue dryness; Chorea. It may be a fatal sign (Deathis predicted within 7 days)
xxiv)	4 ^{1/2} FB	Anasarca or general body swelling; Severe Sunken eyes; It may be a fatal sign (Death is predicted within 9 days)
xxv)	4 ^{1/4} FB	Body tremors; Weakness of both hands and legs; Face becomes blackish in nature; Extreme difficulty in breathing; It may be a fatal sign (Death may be predicted within 2 days)
xxvi)	4 FB	Emaciation of face with diminished eye function; Swelling of both extremities; It may be a fatal sign (Death may be predicted within 5 days)

^{*}FB — Finger-breadth

ANNEX B

(Clauses $\frac{7}{2}$ and $\frac{9.1}{2}$)

MODEL CASE SHEET PROFORMA — CASE REPORTS AND DIAGNOSTIC ACCURACY STUDIES

UHID				No	ame (of the	patie	nt			Оссир	ation	Age	Sex	D_i	iet	Nilam (Tick Appropriately)							
															Veg	NV	Kuriñci	Mullai	Ма	rutam	Neyt	al	Pāla	ıi
Date	Nā	ți kā	lam				Nāṭi naṭai (Pulse Reading) Please Tick Yākkai ilakkaṇam (Body Constitution) Please Ti							ick										
	V	P	K	V	P	K	VP	VK	PV	PK	KV	KP	MI	ζ	V	P	K	VP	VK	PV	PK	KV	KP	MK
WC in FB							Nīrku <u>r</u> i Neykku <u>r</u> i									1								
	Result in Finger Breadth (4-11) WC/1 FB																							
4 FB											Po	<u>l</u> utu ('.	Гіск Аррг	opriately	7)									
1 FB	FB					Kārkālam Kūtirkālam				!	М	u <u>n</u> pa <u>n</u>	ikkālar	lam Pi <u>n</u> pa <u>n</u> ikkālam				Iļavē <u>n</u> i <u>r</u> kālam		Mutuvē <u>n</u> i <u>r</u> kālam				
Clinica	Clinical Findings and Presenting complaints Observations from Manikh							Maṇikkaṭai Reading by Finger Breadth (FBU)																
1																								
3								3																
4									4															
5										5														
Comor DM/HT			Γhyro	oid/He	eart o	liseas	se/Oth	ers (Sp	ecify)				1											
1	mination of Cāram Cennīr						r Ū <u>n</u>				K	Ко <u>l</u> ирри		Ко <u>І</u> ирри І		Е <u>п</u> ри	при 1		Mūļai			Cukkilam/ Cur`ōṇitam		
	prim stitu	-)	S	I	D	S	I	D	S	I	D	S	I	D	S	I	D	S	I	D	S	I	D
S	Subje	ct									~ .				riptio							Tick		
	tegor Readi			 a) True Positive symptoms (TP) – Clinical presentations coinciding with <i>Maṇikkaṭai</i> Reading b) Sequence of True Positive symptoms (SqTP) – Clinical presentations coinciding with multiple Finger breadth readings showing a sequence of origin and progress 																				

IS 19055: 2024

	c) False-positive symptoms (FP) – Subjects not presenting, expressing, or having the symptoms even though the features are shown in the particular reading	
	d) False-negative symptoms (FN) – Subject is presenting the symptom or disease that has no description in the obtained reading	
	e) Investigations: Serological/others (Specify)	
Attachments (If	f) Imaging: X-ray/Scan/others (Specify)	
available)	g) Previous case/Medical record (Copy)	
	h) Photographs/Images of Nīr/Neikuri pattern and its reports	

UHID: Unique Health Identification Number, V: *Vātam*, P: *Pittam*, K: *Kapam*, VP: *Vātapittam*, VK: *Vātakapam*, PV: *Pittavātam*, PK: *Pittakapam*, KV: *Kapavātam*, KP: *Kapapittam*, MK: *Mukkurram*, WC: Wrist Circumference, FB: Finger Breadth, S: Sufficient, I: Increased, D: Decreased.

NOTES

- 1 Manikkațai nūl test results should not be interpreted alone; it must be correlated with clinical information and be interpreted by Registered Siddha Medical Practitioners only to determine final diagnosis.
- 2 Maṇikkaṭai nūl test results may vary based on time of testing, season, physiological condition of the patient, current medication, or nutritional and dietary changes. Please consult your Siddha doctor for any clarification.
- 3 Maņikkaṭai nūl test results cannot be used for Medico-legal purposes.

ANNEX C

(<u>Foreword</u>)

COMMITTEE COMPOSITION

Siddha Sectional Committee, AYD 05

Organization	Representative(s)
National Institute of Siddha, Chennai	Prof Dr R. Meenakumari (<i>Chairperson</i>)
Central Council for Research in Siddha, Chennai	DR SHYAMALA RAJKUMAR DR K SAMRAJ (<i>Alternate</i> I) DR V. AARTHI (<i>Alternate</i> II)
Central Drugs Standard Control Organisation, New Delhi	SHRI SUSHANT SHARMA DR RACHNA PALIWAL (Alternate)
CSIR-Central Institute of Medicinal and Aromatic Plants (CSIR-CIMAP), Lucknow	DR KARUNA SHANKER DR DEBABRATA CHANDA (<i>Alternate</i> I) DR KAPIL DEV (<i>Alternate</i> II)
CSIR-Traditional Knowledge Digital Library Unit (CSIR-TKDL), New Delhi	DR VIJAYALAKSHMI ASTHANA
Department of Pharmacy, Siddha Central Research Institute (CCRS-SCRI), Chennai	DR MS SHREE DEVI DR S. VINAYAK (Alternate)
Drug Testing Laboratory for Testing of A & S, (CCRAS – CSMCARI), Chennai	DR R. ILAVARASAN DR S. CHITRA (Alternate I) MS MONIKA. N (Alternate II)
Government Siddha Medical College, Chennai	Dr D. Sasikumar Dr K. Nalina Saraswathi (<i>Alternate</i>)
Government Siddha Medical College, Palayamkottai	DR T. KOMALAVALLI ALIAS MEHALA DR A. BALAMURUGAN (Alternate I) DR S. SIVANANDAN (Alternate II)
Kokila Siddha Hospital and Research Centre, Madurai	DR J. JEYAVENKATESH DR P. PAVITHRA (Alternate)
National Commission for Indian System of Medicine, New Delhi	DR M.V. MAHADEVAN DR N. J. MUTHUKUMAR (Alternate I) DR V. JASMINE RANJANA (Alternate II)
National Institute of Siddha, Chennai	DR S.SUDHA REVATHY DR V. SUBA (Alternate I) DR B. ANBARASAN (Alternate II)
National Medicinal Plants Board, New Delhi	Dr R. Murugeswaran
Pharmacopoeia Commission for Indian Medicine & Homoeopathy, (PCIM&H), Ghaziabad	DR V. VIJAYAKUMAR DR NIKHIL M. JIRANKALGIKAR (<i>Alternate</i>)
State Drug Licensing Authority (Indian Medicine), Chennai	DR Y.R. MANEKSHAH DR G. SUBASH CHANDRAN (Alternate)
Tamil Nadu Ayurveda, Siddha Unani Manufacturers Association (TASUDMA), Chennai	DR L. SIVAKUMAR DR S. ILANGO (<i>Alternate</i> I) DR J. DINESH KUMAR (<i>Alternate</i> II)

IS 19055: 2024

Organization

Representative(s)

Tamil Nadu Medicinal Plant Farms and Herbal Medicine Corporation Limited (TAMPCOL),

Chennai

The Indian Medical Practitioners' Co-operative Pharmacy & Stores (IMPCOPS), Chennai

DR A. JAHEER HUSSAIN

DR R. KANNAN

DR K. KADER MOHIDEEN (Alternate I) DR S. SUDARSHAN (Alternate II)

The Tamil Nadu Dr MGR Medical University,

Chennai

BIS Directorate General

DR N. KABILAN DR E. M. MANIKGANTAN (Alternate)

SHRIMATI RACHNA SEHGAL, SCIENTIST 'G' AND [REPRESENTING HEAD (AYUSH) DIRECTOR

GENERAL (*Ex-officio*)]

Member Secretary DR KIRUTHIGA G. SCIENTIST 'B'/ASSISTANT DIRECTOR (AYUSH), BIS

Panel for Maņikkaṭai nūl (Wrist Circummetric Sign) Analysis — Code of Practice AYD 05/Panel 02

Organization Representative(s)

Kokila Siddha Hospital and Research Centre, Madurai DR J. JEYAVENKATESH (Convener)

Excel Siddha Medical College, Namakkal DR P. SARAVANA PANDIAN

Kokila Siddha Hospital and Research Centre, Madurai DR P. PAVITHRA

Kokila Siddha Hospital and Research Centre, Madurai Dr S. K. Subhalakshmi This Pade has been Intentionally left blank

This Pade has been Intentionally left blank

Bureau of Indian Standards

BIS is a statutory institution established under the *Bureau of Indian Standards Act*, 2016 to promote harmonious development of the activities of standardization, marking and quality certification of goods and attending to connected matters in the country.

Copyright

BIS has the copyright of all its publications. No part of these publications may be reproduced in any form without the prior permission in writing of BIS. This does not preclude the free use, in the course of implementing the standard, of necessary details, such as symbols and sizes, type or grade designations. Enquiries relating to copyright be addressed to the Head (Publication & Sales), BIS.

Review of Indian Standards

Amendments are issued to standards as the need arises on the basis of comments. Standards are also reviewed periodically; a standard along with amendments is reaffirmed when such review indicates that no changes are needed; if the review indicates that changes are needed, it is taken up for revision. Users of Indian Standards should ascertain that they are in possession of the latest amendments or edition by referring to the website-www.bis.gov.in or www.standardsbis.in.

This Indian Standard has been developed from Doc No.: AYD 05 (23899).

Amendments Issued Since Publication

Amend No.	Date of Issue	Text Affected	

BUREAU OF INDIAN STANDARDS

Headquarters:

Manak Bhavan, 9 Bahadur Shah Zafar Marg, New Delhi 110002

Telephones: 2323 0131, 2323 3375, 2323 9402 Website: www.bis.gov.in

Regional Offices:							
Central : 601/A, Konnectus Tower -1, 6 th Floor, DMRC Building, Bhavbhuti Marg, New Delhi 110002	{ 2323 7617						
Eastern : 8 th Floor, Plot No 7/7 & 7/8, CP Block, Sector V, Salt Lake, Kolkata, West Bengal 700091	2367 0012 2320 9474						
Northern: Plot No. 4-A, Sector 27-B, Madhya Marg, Chandigarh 160019	{ 265 9930						
Southern : C.I.T. Campus, IV Cross Road, Taramani, Chennai 600113	2254 1442 2254 1216						
Western: 5 th Floor/MTNL CETTM, Technology Street, Hiranandani Gardens, Powai Mumbai 400076	25700030 25702715						

Branches: AHMEDABAD, BENGALURU, BHOPAL, BHUBANESHWAR, CHANDIGARH, CHENNAI, COIMBATORE, DEHRADUN, DELHI, FARIDABAD, GHAZIABAD, GUWAHATI, HARYANA (CHANDIGARH), HUBLI, HYDERABAD, JAIPUR, JAMMU, JAMSHEDPUR, KOCHI, KOLKATA, LUCKNOW, MADURAI, MUMBAI, NAGPUR, NOIDA, PARWANOO, PATNA, PUNE, RAIPUR, RAJKOT, SURAT, VIJAYAWADA.