

पारम्परिक औषधियों में उपयोग हेतु
हेमिडेसमस इंडिकस (एल.) आर. ब्र.
जड़ — विशिष्टि

***Hemidesmus indicus* (L.) R. Br Root
for Use in Traditional Medicine —
Specification**

ICS 11.120.10

© BIS 2024



भारतीय मानक ब्यूरो
BUREAU OF INDIAN STANDARDS
मानक भवन, 9 बहादुर शाह ज़फर मार्ग, नई दिल्ली - 110002
MANAK BHAVAN, 9 BAHADUR SHAH ZAFAR MARG
NEW DELHI - 110002

www.bis.gov.in www.standardsbis.in

FOREWORD

This Indian Standard was adopted by Bureau of Indian Standards, after the draft finalized by the Siddha Sectional Committee had been approved by the Ayush Division Council.

Hemidesmus indicus (L.) R. Br. (Fam. Asclepiadaceae), a prostrate or semi-erect laticiferous herb, found throughout India from upper Gangetic plains east-wards to Assam, throughout Central, Western and Southern India up to an elevation of 600 meter. Roots of the plant either dried or fresh are used as an ingredient formulation in traditional Indian system of medicine.

Hemidesmus indicus (L.) R. Br. is also called by various regional names in India. Some of them *Vaga sariva* (Assamese); *Anantamul*, *Shvetashariva* (Bengali); Indian Sarasaparilla (English); *Upalsari*, *Kabri* (Gujarati); *Anantamul* (Hindi); *Namada veru*, *Bili namadaberu*, *Anantamool*, *Sogadeberu*, *Namadaberu* (Kannada); *Nannari*, *Nannar*, *Naruneendi* (Malayalam); *Upalsari*, *Anantamula* (Marathi); *Dralashvan lai*, *Anantamool* (Oriya); *Anantmool*, *Ushbah* (Punjab); *Ananta*, *Gopasuta*, *Sveta Sariva* (Sanskrit); *Ankari muli*, *Cariyam*, *Kamavalli*, *Krishnavalli*, *Nannari*, *Patala muli* (Tamil); *Sugandhi Pala*, *Tella Sugandhi* (Telugu); and *Ushba Hindi* (Urdu).

Other botanical species such as *Ichnocarpus frutescens* (L.) W.T. Aiton, *Decalepis hamiltonii* Wight & Arn. and *Cryptolepis buchananii* Roem. & Schult. are also available in the market under the trade name of *Sariva*.

The standard is one of the series of standards being brought out on ingredients used in formulations of traditional medicine for the advantage of researchers, academicians, students, clinical practitioners, drug manufacturers and raw drug exporters.

In the formulation of this standard, significant assistance has been derived from the Siddha Pharmacopoeia of India, Part I, Vol I, 2008 and Ayurveda Pharmacopoeia of India, Part I, Vol I, 2001 published by the Ministry of Ayush, Government of India. Inputs have also been derived from the information available in the public domain in print and electronic media including authoritative books.

In the formulation of this standard due consideration has been given to the provisions of *The Drugs and Cosmetics Act*, 1940 and Rules framed thereunder. However, this standard is subject to the restrictions imposed under these Rules and Regulations, wherever applicable.

The composition of the Committee responsible for the formulation of this standard is given in [Annex C](#).

For the purpose of deciding whether a particular requirement of this standard is complied with the final value, observed or calculated, expressing the result of a test or analysis, shall be rounded off in accordance with IS 2 : 2022 'Rules for rounding off numeral values (*second revision*)'. The number of significant places retained in the rounded off value should be the same as that of the specified value in the relevant referred standard.

Indian Standard
**HEMIDESMUS INDICUS (L.) R. Br. ROOT FOR USE IN
 TRADITIONAL MEDICINE — SPECIFICATION**

1 SCOPE

This standard prescribes the requirements and method of test for root of *Hemidesmus indicus* (L.) R. Br. (Fam. Asclepiadaceae).

2 REFERENCES

The standards listed in [Annex A](#) contain provisions which through reference in this text, constitute provisions of this standard. At the time of publication, the editions indicated were valid. All standards are subject to revision and parties to agreements based on this standard are encouraged to investigate the possibility of applying the most recent edition of these standards.

3 REQUIREMENTS**3.1 Description****3.1.1 Macroscopic Examination of *Hemidesmus indicus* (L.) R. Br. Root**

Roots are pieces about 30 cm long and 3 mm to 8 mm in diameter, cylindrical, thick, hard, somewhat tortuous, sparsely branched with a few thick rootlets and secondary roots. The roots are dark brown in colour, sometimes with violet-grey tinge, yellow in the center, woody, surrounded by a mealy white cortical layer. The bark is brownish, corky, marked with transverse cracks, longitudinal fissures and is easily detachable from the hard central core. The odour is characteristic and the taste is sweetish, slightly acrid and aromatic.

3.1.2 Microscopic Examination of *Hemidesmus indicus* (L.) R. Br. Root

Transverse section of root shows periderm consisting of three layers of tissues, cork, cork cambium and secondary cortex. The cork cells are radially flattened and rectangular in appearance filled with dark brown contents that gives reactions of tannins. The cork cambium is 2 or 3 layered, compressed and is filled with deep brown contents. The secondary cortex is 3 or 4 layered with cells similar to cork cells. It has very little or no dark brown contents. The secondary phloem consists of sieve elements, parenchyma, phloem ray cells along with several laticiferous ducts. The parenchyma cells are filled with starch grains, measuring diameter 7µm to 10 µm in diameter with

occasional prismatic crystals of calcium oxalate. The laticiferous ducts are scattered in parenchymatous tissue. The cambium is very narrow and the xylem is traversed by narrow medullary rays. The vessels and tracheids are characterized by the presence of pitted markings. Pith is absent and the central region is occupied by woody tissues.

3.1.3 Powder Microscopy of *Hemidesmus indicus* (L.) R. Br. Root

Roots of *Hemidesmus indicus* (L.) R. Br. dried and crushed in the form of fine powder appears brown in colour. The powder shows parenchyma cells filled with oval or rounded starch grains 7 µm to 19 µm in diameter, having 2 to 8 or more components or prismatic calcium oxalate crystals, pieces of laticiferous ducts and vessels with spiral thickenings.

3.2 General

3.2.1 *Hemidesmus indicus* (L.) R. Br. roots shall be free from extraneous/artificial flavours.

3.2.2 *Hemidesmus indicus* (L.) R. Br. roots shall comply with physical, chemical and microbiological requirements given in [Table 1](#).

3.2.3 If any pesticides other than those for which minimum requirements are given in [Table 1](#) are applied to the herb before or after harvesting, those should also be tested. Their limit shall be calculated using the following formula:

$$\frac{ADI \times M}{MDD \times 100}$$

where

ADI = acceptable daily intake of pesticide as published by food and agriculture organization-world health organization (FAO-WHO), in milligrams per kilogram of body mass;

M = body mass in kilograms (60 kg); and

MDD = maximum daily dose of the drug, in kilograms.

Table 1 Requirements for *Hemidesmus indicus* (L.) R. Br. Root(Clauses [3.2.2](#), [3.2.3](#) and [4.2](#))

SI No.	Characteristic	Requirement	Method of Test, Ref to
(1)	(2)	(3)	(4)
i)	Foreign matter, percent by mass, <i>Max</i>	02	6.2 of IS 4333 (Part 1)
ii)	Total ash, percent by mass, <i>Max</i>	04	6 of IS 1797
iii)	Acid insoluble ash, percent by mass, <i>Max</i>	0.5	8 of IS 1797
iv)	Alcohol-soluble Extractive, percent by mass, <i>Min</i>	15	10 of IS 1797
v)	Water soluble Extractive, percent by mass, <i>Min</i>	13	11 of IS 1797
vi)	Aflatoxin (B ₁), ng/kg, <i>Max</i>	2.0	IS 16287
vii)	Aflatoxin (B ₁ + B ₂ + G ₁ + G ₂), ng/kg, <i>Max</i>	5.0	IS 16287
viii)	Lead, ppm, <i>Max</i>	10.0	IS 16913
ix)	Mercury, ppm, <i>Max</i>	1.0	IS 16913
x)	Cadmium, ppm, <i>Max</i>	0.3	IS 16913
xi)	Arsenic, ppm, <i>Max</i>	3.0	
xii)	<i>Staphylococcus aureus</i> , CFU per g	Absent for extract and powder	Annex B
		Absent for plant material	
xii)	<i>Pseudomonas aeruginosa</i> , CFU per g	Absent for extract and powder	Annex B
		Absent for plant material	
xiii)	<i>Salmonella</i> spp., CFU per g	Absent	Annex B
xiv)	<i>Escherichia coli</i> , CFU per g	Absent for plant material	Annex B
		10 for plant material	
xv)	Total microbial plate count, CFU per g, <i>Max</i>	10 ⁵ for extract and powder	Annex B
		10 ⁷ for plant material	
xvi)	Total yeast and mould, CFU per g, <i>Max</i>	10 ³ for extract and powder	Annex B
		10 ⁵ for plant material	
xvii)	Pesticide residues		
a)	Alachlor, (mg/kg), <i>Max</i>	0.02	IS 17924
b)	Aldrin and Dieldrin (sum of), (mg/kg), <i>Max</i>	0.05	
c)	Azinphos-methyl, (mg/kg), <i>Max</i>	1.0	
d)	Bromopropylate, (mg/kg), <i>Max</i>	3.0	
e)	Chlordane (sum of cis-, trans - and Oxythlordane), (mg/kg), <i>Max</i>	0.05	
f)	Chlorfenvinphos, (mg/kg), <i>Max</i>	0.5	
g)	Chlorpyrifos, (mg/kg), <i>Max</i>	0.2	
h)	Chlorpyrifos-methyl, (mg/kg), <i>Max</i>	0.1	
j)	Cypermethrin (and isomers), (mg/kg), <i>Max</i>	1.0	
k)	DDT (sum of p,p'-DDT, o,p'-DDT, p,p'-DDE and p,p'-TDE), (mg/kg), <i>Max</i>	1.0	
m)	Deltamethrin, (mg/kg), <i>Max</i>	0.5	
n)	Diazinon, (mg/kg), <i>Max</i>	0.5	

SI No.	Characteristic	Requirement	Method of Test, Ref to
(1)	(2)	(3)	(4)
p)	Dichlorvos(mg/kg) <i>Max</i>	1.0	IS 17924
q)	Dithiocarbamates (as CS ₂) (mg/kg) <i>Max</i>	2.0	
r)	Endosulfan (sum of isomers and endosulfansulphate) (mg/kg) <i>Max</i>	3.0	
s)	Endrin(mg/kg) <i>Max</i>	0.05	
t)	Ethion(mg/kg) <i>Max</i>	2.0	
u)	Fenitrothion(mg/kg) <i>Max</i>	0.5	
w)	Fenvalerate(mg/kg) <i>Max</i>	1.5	
y)	Fonofos(mg/kg) <i>Max</i>	0.05	
z)	Heptachlor (sum of heptachlor and heptachlor epoxide) (mg/kg) <i>Max</i>	0.05	
aa)	Hexachlorobenzene(mg/kg) <i>Max</i>	0.1	
bb)	Hexachlorocyclohexane isomers (other than γ) (mg/kg) <i>Max</i>	0.3	
cc)	Lindane (γ -hexachlorocyclohexane) (mg/kg) <i>Max</i>	0.6	
dd)	Malathion(mg/kg) <i>Max</i>	1.0	
ee)	Methidathion(mg/kg) <i>Max</i>	0.2	
ff)	Parathion(mg/kg) <i>Max</i>	0.5	
gg)	Parathion-methyl(mg/kg) <i>Max</i>	0.2	
hh)	Permethrin(mg/kg) <i>Max</i>	1.0	
jj)	Phosalone(mg/kg) <i>Max</i>	0.1	
kk)	Piperonylbutoxide(mg/kg) <i>Max</i>	3.0	
mm)	Pirimiphos-methyl(mg/kg) <i>Max</i>	4.0	
nn)	Pyrethrins (sum of) (mg/kg) <i>Max</i>	3.0	
pp)	Quintozene (sum of quintozene, pentachloroaniline and methyl pentachlorophenyl sulphide) (mg/kg) <i>Max</i>	1.0	

NOTES

- 1 Reagents including pure chemicals used shall be of analytical grade.
- 2 Reagent grade water for laboratory use shall be as per IS 1070.
- 3 Pure chemicals shall mean chemicals that do not contain impurities which effect the results of analysis.

4 SAMPLING

4.1 Representative samples of the material shall be drawn and tested for conformity to this specification as prescribed in IS 13145.

4.2 The samples of *Hemidesmus indicus* (L.) R. Br. root shall be tested for ascertaining conformity of the material to the requirements in accordance with the relevant clauses given in col (4) of [Table 1](#).

5 PACKING, STORAGE AND MARKING

5.1 Packing

Hemidesmus indicus (L.) R. Br. roots shall be packed in clean, sound and dry container made up of metal, glass, food grade polymers, wood or jute bags. The wooden boxes or jute bags shall be suitably lined with moisture proof lining which does not impart any foreign smell to the product.

The packing material shall be free from any fungal or insect infestation and should not impart any foreign smell. Each container shall be securely closed and sealed.

5.2 Storage

Hemidesmus indicus (L.) R. Br. roots shall be stored under conditions that prevent contamination and, as far as possible, deterioration. Storage area shall be clean, well-ventilated, protected from direct light, moisture and insects and rodents.

5.3 Marking

The following particulars shall be legibly and indelibly marked or labelled on each pack:

- a) Name of the material including part of the plant, botanical name, and trade name or brand name, if any;

IS 19090 : 2024

- b) Name and address of the producer or packer;
- c) Batch number;
- d) Net quantity;
- e) State and country of production;
- f) Date of packing (MM/YYYY);
- g) Instructions for storage; and
- h) Any other information requested by the buyer, such as the date of harvesting (MM/YYYY) (if known).

The above information, or part of it, may instead appear in the documentation after agreement between the buyer and the seller.

5.4 BIS Certification Marking

The product(s) conforming to the requirements of this standard may be certified as per the conformity assessment schemes under the provisions of the *Bureau of Indian Standards Act, 2016* and the Rules and Regulations framed there under, and the product(s) may be marked with the Standard Mark.

ANNEX A

(Clause 2)

LIST OF STANDARDS REFERRED

<i>IS No.</i>	<i>Title</i>	<i>IS No.</i>	<i>Title</i>
IS 1070 : 2023	Reagent grade water — Specification (<i>fourth revision</i>)		G ₁ and G ₂ in cereals, nuts and derived products — High performance liquid chromatographic method
IS 1797 : 2017	Spices and condiments — Methods of test (<i>third revision</i>)	IS 16913 : 2018	Methods of test for cosmetics — Determination of heavy metals (arsenic, cadmium, lead and mercury) by atomic absorption spectrometry (AAS)
IS 4333 : 2018	Methods of analysis for food grains: Part 1 Refractions (<i>third revision</i>)		
IS 13145 : 2014	Spices and condiments — Methods of sampling (<i>second revision</i>)	IS 17924 : 2022	Determination of pesticide residue in herbal materials
IS 16287 : 2015/ ISO 16050 : 2003	Foodstuffs — Determination of aflatoxin B ₁ , and the total content of aflatoxins B ₁ , B ₂ ,		

To access Indian Standards click on the link below:

https://www.services.bis.gov.in/php/BIS_2.0/bisconnect/knowyourstandards/Indian_standards/isdetails/

ANNEX B

[Table 1, Sl No. (xiii), (xiv), (xv), (xvi), (xvii) and (xviii)]

TESTS FOR MICROBIAL LIMITS

B-1 GENERAL

The tests are designed for the estimation of the number of viable aerobic microorganisms present and for detecting the presence of designated microbial species in the extract. The term 'growth' is used to designate the presence and presumed proliferation of viable microorganisms.

B-2 APPARATUS**B-2.1 Oven for Dry Sterilization****B-2.2 Autoclave for Wet Sterilization****B-2.3 Incubator****B-2.4 Water Bath****B-2.5 pH Meter**

B-2.6 Sterile Membrane Filters — 50 mm in diameter

B-2.7 Colony Counting Equipment**B-2.8 Analytical Balance****B-2.9 Standard Glassware****B-3 MEDIA**

Culture media may be prepared as given below or dehydrated culture media may be used provided that, when reconstituted as directed by the manufacturer, they have similar ingredients and/or yield media comparable to those obtained from the formulae given below.

Where agar is specified in a formula, use agar that has a moisture content of not more than 15 percent. Where water is called for in a formula, use purified water. Unless otherwise indicated, the media should be sterilized by heating in an autoclave (15 psi) at 121 °C for 15 min. In preparing media by the formulas given below, dissolve the soluble solids in the water, using heat, if necessary, to effect complete solution, add solutions of 0.1 N hydrochloric acid or 0.1 N sodium hydroxide in quantities sufficient to yield the required pH in the medium when it is ready for use. Determine the pH at 25 °C ± 2 °C.

B-3.1 Baird Parker Agar Medium**B-3.1.1 Composition**

Pancreatic digest of casein 10.0 g

Beef extract	5.0 g
Yeast extract	1.0 g
Lithium chloride	5.0 g
Agar	20.0 g
L-Glycine	12.0 g
Sodium pyruvate	10.0 g
Water	1 000 ml

B-3.1.2 Preparation

Suspend the components in 1 000 ml of water, heat with frequent agitation and boil for 1 min. Sterilize, cool in between 45 °C to 50 °C, add 10 ml of a one percent w/v solution of sterile potassium tellurite and 50 ml of egg yolk emulsion. Mix thoroughly, but gently and pour into plates. (Prepare the egg yolk emulsion by disinfecting the surface of whole shell eggs, aseptically cracking the eggs, and separating out intact yolks into a sterile graduated cylinder. Add sterile saline solution, get a 3 : 7 of egg yolk to saline. Add to a sterile blender cup and mix at high speed for 5 s). Adjust the pH after sterilization to 6.8 ± 0.2.

B-3.2 Bismuth Sulphite Agar Medium**B-3.2.1 Composition**

Solution (1)

Beef extract	6 g
Peptone	10 g
Agar	24 g
Ferric citrate	0.4 g
Brilliant green	10 mg
Water	1 000 ml

Solution (2)

Ammonium bismuth citrate	3 g
Sodium sulphite	10 g
Anhydrous disodium hydrogen phosphate	5 g
Dextrose monohydrate	5 g
Water	100 ml

B-3.2.2 Preparation

Suspend the components of Solution (1) in 1 000 ml of water. Heat to boiling to dissolve the medium completely. Sterilize by maintaining at 115 °C for 30 min.

Suspend the components of Solution (2) in 100 ml of water. Heat to boiling to dissolve the medium completely. Do not autoclave.

Add 1 volume of Solution (2) to 10 volumes of Solution (1) which was previously melted and cooled to a temperature of 55 °C. Bismuth sulphite agar medium should be stored at 2 °C to 8 °C for 5 days before use.

B-3.3 Brilliant Green Agar Medium

B-3.3.1 Composition

Peptone	10.0 g
Yeast extract	3.0 g
Lactose	10.0 g
Sucrose	10.0 g
Sodium chloride	5.0 g
Phenol red	80.0 g
Brilliant green	12.5 mg
Agar	12.0 g
Water	1 000 ml

B-3.3.2 Preparation

Mix the components, allow to stand for 15 min, sterilize by maintaining at 115 °C for 30 min and mix before pouring.

B-3.4 Buffered Sodium Chloride Peptone Solution pH 7.0

B-3.4.1 Composition

Potassium dihydrogen phosphate	3.56 g
Disodium hydrogen phosphate dihydrate	7.23 g
Sodium chloride	4.30 g
Peptone (meat or casein)	1.0 g
Water	1 000 ml

B-3.4.2 Preparation

Mix the components and heat if necessary to dissolve the medium completely. 0.1 percent to 1.0 percent *w/v* polysorbate 20 or polysorbate 80 may be added. Sterilize by heating in an autoclave at 121 °C for 15 min.

B-3.5 Casein Soyabean Digest Agar Medium

B-3.5.1 Composition

Pancreatic digest of casein	15.0 g
Papaic digest of soyabean meal	5.0 g
Sodium chloride	5.0 g
Agar	15.0 g
Water	1 000 ml

B-3.5.2 Preparation

Suspend the components in water. Heat to boiling to dissolve the medium completely. Sterilize at 121 °C for 15 min in an autoclave and adjust the pH after sterilization to 7.3 ± 0.2.

B-3.6 Cetrinide Agar Medium

B-3.6.1 Composition

Pancreatic digest of gelatin	20.0 g
Magnesium chloride	1.4 g
Potassium sulphate	10.0 g
Cetrinide	0.3 g
Agar	13.6 g
Glycerin	10.0 g
Water	1 000 ml

B-3.6.2 Preparation

Dissolve the components in 1 000 ml of water. Heat to boiling for 1 min with shaking. Sterilize at 121 °C for 15 min in an autoclave and adjust the pH after sterilization to 7.0 to 7.4.

B-3.7 Deoxycholate Citrate Agar Medium

B-3.7.1 Composition

Beef extract	5.0 g
Peptone	5.0 g
Lactose	10.0 g
Trisodium citrate	8.5 g
Sodium thiosulphate	5.4 g
Ferric citrate	1.0 g
Sodium deoxycholate	5.0 g
Neutral red	0.02 g
Agar	12.0 g
Water	1 000 ml

B-3.7.2 Preparation

Mix the components and allow to stand for 15 min. Gently boil with continuous stirring and continue boiling until solution is complete. Cool to 80 °C, mix, pour and cool rapidly.

Care should be taken not to overheat deoxycholate citrate agar during preparation. It should not be re-melted and the surface of the plates should be dried before use.

B-3.8 Fluid Casein Digest Soya Lecithin Polysorbate 20 Medium

B-3.8.1 Composition

Pancreatic digest of casein	20 g
Soya lecithin	5 g
Polysorbate 20	40 ml
Water	1 000 ml

B-3.8.2 Preparation

Dissolve the pancreatic digest of casein and soya lecithin in water, heating in a water bath at 48 °C to 50 °C for about 30 min to effect solution.

IS 19090 : 2024

Add polysorbate 20, mix and dispense as desired. Sterilize at 121 °C for 15 min in an autoclave.

B-3.9 Fluid Lactose Medium

B-3.9.1 Composition

Beef extract	3.0 g
Pancreatic digest of gelatin	5.0 g
Lactose	5.0 g
Water	1 000 ml

B-3.9.2 Preparation

Suspend the components in 1 000 ml water. Heat if necessary to dissolve the medium completely. Sterilize at 121 °C for 15 min in an autoclave. Cool as quickly as possible after sterilization. Adjust the pH after sterilization to 6.9 ± 0.2 .

B-3.10 Lactose Broth Medium

B-3.10.1 Composition

Beef extract	3.0 g
Pancreatic digest of gelatin	5.0 g
Lactose	5.0 g
Water	1 000 ml

B-3.10.2 Preparation

Suspend the components in water and heat if necessary to dissolve the medium completely. Sterilize at 121 °C for 15 min in an autoclave. Adjust the pH after sterilization to 6.9 ± 0.2 .

B-3.11 Levine Eosin Methylene Blue Agar Medium

B-3.11.1 Composition

Pancreatic digest of gelatin	10.0 g
Dibasic potassium phosphate	2.0 g
Agar	15.0 g
Lactose	10.0 g
Eosin Y	400 mg
Methylene blue	65 mg
Water	1 000 ml

B-3.11.2 Preparation

Dissolve the pancreatic digest of gelatin, dibasic potassium phosphate and agar in water with warming and allow to cool. Just prior to use, liquify the gelled agar solution and the remaining ingredients, as solutions, in the following amounts and mix. For each 100 ml of the liquified agar solution use 5 ml of a 20 percent w/v solution of lactose, 2 ml of a 2 percent w/v solution of eosin Y and 2 ml of a 0.33 percent w/v solution of methylene blue. The finished medium may not be

clear. Adjust the pH after sterilization to 7.1 ± 0.2 .

B-3.12 MacConkey Agar Medium

B-3.12.1 Composition

Pancreatic digest of gelatin	17.0 g
Peptone (meat and casein, equal parts)	3.0 g
Lactose	10.0 g
Sodium chloride	5.0 g
Bile salts	1.5 g
Agar	13.5 g
Neutral red	30 mg
Crystal violet	1 mg
Water	1 000 ml

B-3.12.2 Preparation

Boil the mixture of solids and water for 1 min to effect solution. Sterilize at 121 °C for 15 min in an autoclave. Adjust the pH after sterilization to 7.1 ± 0.2 .

B-3.13 MacConkey Broth Medium

B-3.13.1 Composition

Pancreatic digest of gelatin	20.0 g
Lactose	10.0 g
Dehydrated ox bile	5.0 g
Bromocresol purple	10 mg
Water	1 000 ml

B-3.13.2 Preparation

Suspend the components in 1 000 ml of water and heat if necessary to dissolve the medium completely. Sterilize at 121 °C for 15 min in an autoclave and adjust the pH after sterilization to 7.3 ± 0.2 .

B-3.14 Mannitol Salt Agar Medium

B-3.14.1 Composition

Pancreatic digest of gelatin	5.0 g
Peptic digest of animal tissue	5.0 g
Beef extract	1.0 g
D-mannitol	10.0 g
Sodium chloride	75.0 g
Agar	15.0 g
Phenol red	25 mg
Water	1 000 ml

B-3.14.2 Preparation

Mix the components, heat with frequent agitation and boil for 1 min to effect solution. Sterilize at 121 °C for 15 min in an autoclave and adjust the pH after sterilization to 7.4 ± 0.2 .

B-3.15 Nutrient Broth Medium**B-3.15.1 Composition**

Beef extract	10.0 g
Peptone	10.0 g
Sodium chloride	5 mg
Water	1 000 ml

B-3.15.2 Preparation

Dissolve the components with the aid of heat. Adjust the pH to 8.0 to 8.4 with 5 M sodium hydroxide and boil for 10 min. Filter and sterilize by maintaining at 115 °C for 30 min and adjust pH to 7.3 ± 0.1 .

B-3.16 Nutrient Agar Medium

Nutrient broth gelled by the addition of 1 percent to 2 percent *w/v* of agar.

B-3.17 Pseudomonas Agar Medium for Detection of Fluorescein**B-3.17.1 Composition**

Pancreatic digest of casein	10.0 g
Peptic digest of animal tissue	10.0 g
Anhydrous dibasic potassium phosphate	1.5 g
Magnesium sulphate hepta hydrate	1.5 g
Glycerin	10.0 ml
Agar	15.0 g
Water	1 000 ml

B-3.17.2 Preparation

Dissolve the solid components in water before adding glycerin. Heat with frequent agitation and boil for 1 min to effect solution. Sterilize at 121 °C for 15 min in an autoclave and adjust the pH after sterilization to 7.2 ± 0.2 .

B-3.18 Pseudomonas Agar Medium for Detection of Pyocyanin**B-3.18.1 Composition**

Pancreatic digest of gelatin	20.0 g
Anhydrous magnesium chloride	1.4 g
Anhydrous potassium sulphate	10.0 g
Agar	15.0 g
Glycerin	10.0 ml
Water	1 000 ml

B-3.18.2 Preparation

Dissolve the solid components in water before adding glycerin. Heat with frequent agitation and boil for 1 min to effect solution. Sterilize at 121 °C for 15 min in an autoclave and adjust the pH after sterilization to 7.2 ± 0.2 .

B-3.19 Sabouraud Dextrose Agar Medium**B-3.19.1 Composition**

Dextrose	40 g
Peptic digest of animal tissue and pancreatic digest of casein (1 : 1)	10 g
Agar	15 g
Water	1 000 ml

B-3.19.2 Preparation

Mix the components and heat to boiling to dissolve completely. Sterilize at 121 °C for 15 min in an autoclave and adjust the pH after sterilization to 5.6 ± 0.2 .

B-3.20 Sabouraud Dextrose Agar Medium with Antibiotics

To 1 000 ml of Sabouraud dextrose agar medium, add 0.1 g of benzyl penicillin sodium and 0.1 g of tetracycline HCL or alternatively add 50 mg of chloramphenicol immediately before use.

B-3.21 Selenite F Broth**B-3.21.1 Composition**

Peptone	5 g
Lactose	4 g
Disodium hydrogen phosphate	10 g
Sodium hydrogen selenite	4 g
Water	1 000 ml

B-3.21.2 Preparation

Suspend the components in water and mix well. Warm to dissolve the medium completely. Distribute in sterile containers and sterilize by maintaining at 100 °C for 30 min.

B-3.22 Fluid Selenite Cystine Medium**B-3.22.1 Composition**

Pancreatic digest of casein	5.0 g
Lactose	4.0 g
Disodium phosphate	10.0 g
Sodium hydrogen selenite	4.0 g
L-Cystine	10.0 mg
Water	1 000 ml

B-3.22.2 Preparation

Suspend the components in water and heat in flowing steam for 15 min. Adjust the final pH to 7.0 ± 0.2 . Do not sterilize.

B-3.23 Tetrathionate Broth Medium**B-3.23.1 Composition**

Beef extract	0.9 g
--------------	-------

IS 19090 : 2024

Peptone	4.5 g
Yeast extract	1.8 g
Sodium chloride	4.5 g
Calcium carbonate	25.0 g
Sodium thiosulphate	40.7 g
Water	1 000 ml

B-3.23.2 Preparation

Dissolve the solids in 1 000 ml of water and heat the solution to boil. On the day of use, add a solution prepared by dissolving 5 g of potassium iodide and 6 g of iodine in 20 ml of water.

B-3.24 Tetrathionate Bile Brilliant Green Broth Medium

B-3.24.1 Composition

Peptone	8.6 g
Dehydrated ox bile	8.0 g
Sodium chloride	6.4 g
Calcium carbonate	20.0 g
Potassium tetrathionate	20.0 g
Brilliant green	70 mg
Water	1 000 ml

B-3.24.2 Preparation

Suspend the components in 1 000 ml of water. Heat just to boiling. Do not autoclave or reheat. Adjust the pH so that after heating it is 7.0 ± 0.2 .

B-3.25 Triple Sugar Iron Agar Medium

B-3.25.1 Composition

Beef extract	3.0 g
Yeast extract	3.0 g
Peptone	20.0 g
Lactose	10.0 g
Sucrose	10.0 g
Dextrose monohydrate	1.0 g
Ferrous sulphate	0.2 g
Sodium chloride	5.0 g
Sodium thiosulphate	0.3 g
Phenol red	24 mg
Water	1 000 ml
Agar	12.0 g

B-3.25.2 Preparation

Mix the components in 1 000 ml of water, allow standing for 15 min. Heat to boiling to dissolve the medium completely. Mix well and distribute into test tubes and sterilize by maintaining at 121 °C for 15 min. Allow the medium to stand in a sloped form with a butt about 2.5 cm long.

B-3.26 Urea Broth Medium

B-3.26.1 Composition

Potassium dihydrogen orthophosphate	9.1 g
-------------------------------------	-------

Anhydrous disodium hydrogen phosphate	9.5 g
Urea	20.0 g
Yeast extract	0.1 g
Phenol red	10 mg
Water	1 000 ml

B-3.26.2 Preparation

Mix the components, sterilize by filtration and distribute aseptically in sterile containers.

B-3.27 Vogel Johnson Agar Medium

B-3.27.1 Composition

Pancreatic digest of casein	10.0 g
Yeast extract	5.0 g
Mannitol	10.0 g
Dibasic potassium phosphate	5.0 g
Lithium chloride	5.0 g
Glycine	10.0 g
Agar	16.0 g
Phenol red	25.0 mg
Water	1 000 ml

B-3.27.2 Preparation

Suspend the components in 1000 ml of water. Boil the solution of solids for 1 min. Sterilize at 121 °C for 15 min in an autoclave. Cool to 45 °C to 50 °C and add 20 ml of 1 percent w/v sterile solution of potassium tellurite. Adjust the pH after sterilization to 7.0 ± 0.2 .

B-3.28 Xylose Lysine Deoxycholate Agar Medium

B-3.28.1 Composition

Xylose	3.5 g
l-Lysine	5.0 g
Lactose	7.5 g
Sucrose	7.5 g
Sodium chloride	5.0 g
Yeast extract	3.0 g
Phenol red	80 mg
Agar	13.5 g
Sodium deoxycholate	2.5 g
Sodium thiosulphate	6.8 g
Ferric ammonium citrate	800 mg
Water	1 000 ml

B-3.28.2 Preparation

Suspend the components in 1 000 ml of water. Heat with frequent agitation until the medium boils. Do not overheat or sterilize. Transfer at once to a water bath maintained at about 50 °C and pour into plates as soon as the medium has cooled. Adjust the final pH to 7.4 ± 0.2 .

B-4 SAMPLING

Use 10 ml or 10 g specimens for each of the tests

specified in the Indian Standard.

B-5 PRECAUTIONS

The microbial limit tests should be carried out under conditions designed to avoid accidental contamination during the test. The precautions taken to avoid contamination must be such that they do not adversely affect any microorganisms that should be revealed in the test.

B-6 PRELIMINARY TESTING

Prior to doing the tests, inoculate diluted extracts being examined with separate viable cultures of *Escherichia coli*, *Salmonella* species, *Pseudomonas aeruginosa* and *Staphylococcus aureus*. This is done by adding 1 ml of 24 h broth culture containing not less than 1 000 microorganisms to the first dilution (in buffer solution pH 7.2, fluid soyabean casein digest medium or fluid lactose medium) of the test material and following the test procedure. If the organisms fail to grow in the relevant medium the procedure should be modified by (a) increasing the volume of diluent with the quantity of test material remaining the same, or (b) incorporating a sufficient quantity of a suitable inactivating agent in the diluents, or (c) combining the afore mentioned modifications so as to permit growth of the organisms in the media. If inhibitory substances are present in the extracts, 0.5 percent of soya lecithin and 4 percent of polysorbate 20 may be added to the culture medium. Alternatively, repeat the test as described in the previous paragraph, using fluid casein digest soya lecithin polysorbate media, to demonstrate neutralization of preservatives or other antimicrobial agents in the test material. Where inhibitory substances are contained in the extracts and the latter is soluble, the membrane filtration method described under total aerobic microbial count may be used. If in spite of incorporation of suitable inactivating agents and a substantial increase in the volume of diluent it is still not possible to recover the viable cultures described above and where the article is not suitable for applying the membrane filtration method, it can be assumed that the failure to isolate the inoculated organism may be due to the bactericidal activity of the product. This may indicate that the article is not likely to be contaminated with the given species of microorganisms. However, monitoring should be continued to establish the spectrum of inhibition and bactericidal activity of the article.

B-7 TOTAL AEROBIC MICROBIAL COUNT

Pre-treat the extracts and raw materials being examined as described below.

NOTE — The raw materials need to be ground as a coarse powder before analysis.

B-7.1 Water Soluble Products

Dissolve 10 g or dilute 10 ml of the extract preparation being examined, unless otherwise specified, in buffered sodium chloride peptone solution pH 7.0 or any other suitable medium shown to have no antimicrobial activity under the conditions of test and adjust the volume to 100 ml with the same medium. If necessary, adjust the pH to about 7.

B-7.2 Products Insoluble in Water (Non-Fatty)

Suspend 10 g or 10 ml of the extract preparation being examined, unless otherwise specified, in buffered sodium chloride peptone solution pH 7.0 or any other suitable medium shown not to have antimicrobial activity under the conditions of the test and dilute to 100 ml with the same medium. If necessary, divide the preparation being examined and homogenize the suspension mechanically. A suitable surface-active agent such as 0.1 percent w/v of polysorbate 80 may be added to assist the suspension of poorly wettable substances. If necessary, adjust pH of the suspension to about 7.

B-7.3 Fatty Products

Homogenize 10 g or 10 ml of the extract preparation being examined, unless otherwise specified, with 5 g of polysorbate 20 or polysorbate 80. If necessary, heat to not more than 40 °C. Mix carefully while maintaining the temperature in the water bath or in an oven. Add 85 ml of buffered sodium chloride peptone solution pH 7.0 or any other suitable medium shown to have no antimicrobial activity under the conditions of the test, heated to not more than 40 °C if necessary. Maintain this temperature for the shortest time necessary for formation of an emulsion and in any case for not more than 30 min. If necessary, adjust the pH to about 7.

B-7.4 Examination of the Sample

Determine the total aerobic microbial count in the extract being examined by any of the following methods.

B-7.5 Membrane Filtration

Use membrane filters 50 mm in diameter and having a nominal pore size not greater than 0.45 µm the effectiveness of which in retaining bacteria has been established for the type of preparation being examined. Transfer 10 ml or a quantity of each dilution containing 1 g of the preparation being examined to each of two membrane filters and filter immediately. If necessary, dilute the pretreated extract preparation so that a colony count of 10 to 100 may be expected. Wash each membrane by filtering

through it three or more successive quantities, each of about 100 ml, of a suitable liquid such as buffered sodium chloride peptone solution pH 7.0. For fatty substances add to the liquid polysorbate 20 or polysorbate 80. Transfer one of the membrane filters, intended for the enumeration of bacteria, to the surface of a plate of casein soyabean digest agar and the other, intended for the enumeration of fungi, to the surface of a plate of Sabouraud dextrose agar with antibiotics.

Incubate the plates for 5 days, unless a more reliable count is obtained in shorter time, at 30 °C to 35 °C in the test for bacteria and 20 °C to 25 °C in the test for fungi. Count the number of colonies that are formed. Calculate the number of microorganisms per g or per ml of the extract preparation being examined, if necessary, count bacteria and fungi separately.

B-7.6 Plate Count for Bacteria

Using petri dishes 9 cm to 10 cm in diameter, add to each dish a mixture of 1 ml of the pretreated extract preparation and about 15 ml of liquified casein soyabean digest agar at not more than 45 °C.

Alternatively, spread the pretreated extract preparation on the surface of the solidified medium in a petri dish of the same diameter. If necessary, dilute the pretreated extract preparation as described above so that a colony count of not more than 300 may be expected. Prepare at least two such petri dishes using the same dilution and incubate at 30 °C to 35 °C for 5 days, unless a more reliable count is obtained in a shorter time. Count the number of colonies that are formed. Calculate the results using plates with the greatest number of colonies but taking 300 colonies per plate as the maximum consistent with good evaluation.

B-7.7 Plate Count for Fungi

Proceed as described in the test for bacteria but use Sabouraud dextrose agar with antibiotics in place of casein soyabean digest agar and incubate the plates at 20 °C to 25 °C for 5 days, unless a more reliable count is obtained in a shorter time. Calculate the results using plates with not more than 100 colonies.

B-7.8 Multiple Tube or Serial Dilution Method

In each of fourteen test tubes of similar size place 9.0 ml of sterile fluid soyabean casein digest medium. Arrange twelve of the tubes in four sets of three tubes each. Put aside one set of three tubes to serve as controls. Into each of three tubes of one set ("100") and into fourth tube (A) pipette 1 ml of the solution of suspension of the test specimen (extract) and mix. From tube A pipette 1 ml of its contents into the one remaining tube (B) not

included in the set and mix. These two tubes contain 100 mg (or 100 µl) and 10 mg (or 10 µl) of the specimen respectively. Into each of the second set ("10") of three tubes pipette 1 ml from tube A, and into each tube of the third set ("1") pipette 1 ml from tube B. Discard the unused contents of tube A and B. Close well and incubate all of the tubes.

Following the incubation period, examine the tubes for growth. The three control tubes remain clear. Observations in the tubes containing the test specimen, when interpreted by reference to [Table 2](#), indicate the most probable number of microorganisms per g or per ml of the test specimen.

B-8 TESTS FOR SPECIFIED MICROORGANISMS

B-8.1 Pre-Treatment of the Extract Sample Being Examined

Proceed as described under the test for total aerobic microbial count but using lactose broth or any other suitable medium shown to have no antimicrobial activity under the conditions of test in place of buffered sodium chloride peptone solution pH 7.0.

B-8.2 *Escherichia coli*

Place the prescribed quantity in a sterile screw capped container, add 50 ml of nutrient broth, shake, allow to stand for 1 h (4 h for gelatin) and shake again. Loosen the cap and incubate at 37 °C for 18 h to 24 h.

B-8.2.1 Primary Test

Add 1.0 ml of the enrichment culture to a tube containing 5 ml of MacConkey broth. Incubate in a water bath at 36 °C to 38 °C for 48 h. If the contents of the tube show acid and gas, carry out the secondary test.

B-8.2.2 Secondary Test

Add 0.1 ml of the contents of the tubes containing (a) 5 ml of MacConkey broth and (b) 5 ml of peptone water. Incubate in a water bath at 43.5 °C to 44.5 °C for 24 h and examine tube (a) for acid and gas and tube (b) for indole. To test for indole, add 0.5 ml of Kovac's reagent, shake well and allow to stand for 1 min. If a red colour is produced in the reagent layer indole is present. The presence of acid and gas and of indole in the secondary test indicates the presence of *Escherichia coli*.

Carry out a control test by repeating the primary and secondary tests, adding 1.0 ml of the enrichment culture and a volume of broth containing 10 *Escherichia coli* to 50 *Escherichia coli* (NCTC 9002) organisms, prepared from a 24 h culture in nutrient broth, to 5 ml of MacConkey

broth. The test is not valid unless the results indicate that the control contains *Escherichia coli*.

B-8.2.3 Alternative Test

By means of an inoculating loop, streak a portion from the enrichment culture (obtained in the previous test) on the surface of MacConkey agar medium. Cover and invert the dishes and incubate.

Upon examination, if none of the colonies are brick red in colour and have a surrounding zone of precipitated bile the sample meets the requirements of the test for the absence of *Escherichia coli*. If the colonies described above are found, transfer the suspect colonies individually to the surface of Levine eosin methylene blue agar medium, plated on petri dishes. Cover and invert the plates and incubate. Upon examination, if none of the colonies exhibits both a characteristic metallic sheen under reflected light and a blue-black appearance under transmitted light, the sample meets the requirements of the test for the absence of *Escherichia coli*. The presence of *Escherichia coli* may be confirmed by further suitable cultural and biochemical tests.

B-8.3 Salmonella

Transfer a quantity of the pretreated preparation being examined containing 1 g or 1 ml of the product to 100 ml of nutrient broth in a sterile screw capped jar, shake, allow to stand for 4 h and shake again. Loosen the cap and incubate at 35 °C to 37 °C for 24 h.

B-8.3.1 Primary Test

Add 1.0 ml of the enrichment culture to each of the two tubes containing (a) 10 ml of selenite F broth and (b) tetrathionate bile brilliant green broth and incubate at 36 °C to 38 °C for 48 h. From each of these two cultures subculture on at least two of the following four agar media: bismuth sulphate agar, brilliant green agar, deoxycholate citrate agar and xylose lysine deoxycholate agar. Incubate the plates at 36 °C to 38 °C for 18 h to 24 h. Upon examination, if none of the colonies conforms to the description given in Table 3, the sample meets the requirements of the test for the absence of the genus *Salmonella*. If any colonies conforming to the description in Table 3 are produced, carry out the secondary test.

Table 2 Most probable Total Count by Multiple Tube or Serial Dilution Method

(Clause B-7.8)

Sl No.	Observed Combination of Numbers of Tubes Showing Growth in Each Set Number of mg (or ml) of Specimen per Tube			Most Probable Number of Microorganisms per g or per ml
	100 (100 µl)	10 (10 µl)	1 (1 µl)	
(1)	(2)	(3)	(4)	(5)
i)	3	3	3	> 1 100
ii)	3	3	2	1 100
iii)	3	3	1	500
iv)	3	3	0	200
v)	3	2	3	290
vi)	3	2	2	210
vii)	3	2	1	150
viii)	3	2	0	90
ix)	3	1	3	160
x)	3	1	2	120
xi)	3	1	1	70
xii)	3	1	0	40
xiii)	3	0	3	95
xiv)	3	0	2	60
xv)	3	0	1	40
xvi)	3	0	0	23

Table 3 Interpretation of the Confirmatory Tests for *Salmonella*(Clauses [B-8.3.1](#) and [8.3.2](#))

Sl No.	Medium	Description of Colony
(1)	(2)	(3)
i)	Bismuth sulphite agar	Black or green
ii)	Brilliant green agar	Small, transparent and colourless, or opaque, pinkish or white (frequently surrounded by a pink or red zone)
iii)	Deoxycholate citrate agar	Colorless and opaque, with or without black centers
iv)	Xylose lysine deoxycholate agar	Red with or without black centres

B-8.3.2 Secondary Test

Subculture any colonies showing the characteristics given in [Table 3](#) in triple sugar iron agar by first inoculating the surface of the slope and then making a stab culture with the same inoculating needle, and at the same time inoculate a tube of urea broth. Incubate at 36 °C to 38 °C for 18 h to 24 h. The formation of acid and gas in the stab culture (with or without concomitant blackening) and the absence of acidity from the surface growth in the triple sugar iron agar, together with the absence of a red color in the urea broth, indicate the presence of *Salmonella*. If acid but no gas is produced in the stab culture, the identity of the organisms should be confirmed by agglutination tests.

Carry out the control test by repeating the primary and secondary tests using 1.0 ml of the enrichment culture and a volume of broth containing 10 *Salmonella abony* to 50 *Salmonella abony* (NCTC 6017) organisms, prepared from a 24 h culture in nutrient broth, for the inoculation of the tubes (a) and (b). The test is not valid unless the results indicate that the control contains *Salmonella*.

B-8.4 *Pseudomonas aeruginosa*

Pretreat the preparation being examined as described above and inoculate 100 ml of fluid soyabean casein digest medium with a quantity of the solution, suspension or emulsion thus obtained containing 1 g or 1 ml of the preparation being examined. Mix and incubate at 35 °C to 37 °C for 24 h to 48 h. Examine the medium for growth and if growth is present, streak a portion of the medium on the surface of cetrimide agar medium, each plated on petri dishes. Cover and incubate at 35 °C to 37 °C for 18 h to 24 h.

If, upon examination, none of the plates contains colonies having the characteristics listed in [Table 4](#) for the media used, the sample meets the requirement for freedom from *Pseudomonas aeruginosa*. If any colonies conforming to the description in [Table 4](#) are produced, carry out the oxidase and pigment tests. Streak representative suspect colonies from the agar surface of cetrimide agar on the surfaces of *Pseudomonas* agar medium for detection of fluorescein and *Pseudomonas* agar medium for detection of pyocyanin contained in petri dishes. Cover and invert the inoculated media and incubate at 33 °C to 37 °C for not less than 3 days. Examine the streaked surfaces under ultra violet light. Examine the plates to determine whether colonies conforming to the description in [Table 4](#) are present. If growth of suspect colonies occurs, place 2 drops or 3 drops of a freshly prepared 1 percent w/v solution of N,N,N1,N1-tetramethyl-4-phenylenediamine dihydrochloride on filter paper and smear with the colony. If there is no development of a pink color, changing to purple, the sample meets the requirements of the test for the absence of *Pseudomonas aeruginosa*.

B-8.5 *Staphylococcus aureus*

Proceed as described under *Pseudomonas aeruginosa*, if upon examination of the incubated plates, none of them contains colonies having the characteristics listed in the media used, the sample meets the requirements for the absence of *Staphylococcus aureus*. If growth occurs, carry out the coagulase test. Transfer representative suspect colonies from the agar surface of any of the media listed in [Table 5](#) to individual tubes, each containing 0.5 ml of mammalian, preferably rabbit or horse, plasma with or without additives.

Table 4 Interpretation of the Confirmatory Tests for *Pseudomonas aeruginosa*(Clause [B-8.4](#))

SI No.	Selective Medium	Characteristic Colonial Morphology	Fluorescence in UV Light	Oxidase Test	Gram Stain
(1)	(2)	(3)	(4)	(5)	(6)
i)	Cetrimide agar	Generally greenish	Greenish	Positive	Negative rods
ii)	<i>Pseudomonas</i> agar medium for detection of fluorescein	Generally colorless to yellowish	Yellowish	Positive	Negative rods
iii)	<i>Pseudomonas</i> agar for detection of pyocyanin	Generally greenish	Blue	Positive	Negative rods

Table 5 Interpretation of the Confirmatory Tests for *Staphylococcus aureus*(Clause [B-8.5](#))

SI No.	Selective Medium	Characteristic Colonial Morphology	Gram Stain
(1)	(2)	(3)	(4)
i)	Vogel johnson agar	Black surrounded by yellow zones	Positive cocci (in clusters)
ii)	Mannitol salt agar	Yellow colonies with yellow zones	Positive cocci (in clusters)
iii)	Baird parker agar	Black, shiny, surrounded by clear zones of 2 mm to 5 mm	Positive cocci (in clusters)

Incubate in water bath at 37 °C examining the tubes at 3 h and subsequently at suitable intervals up to 24 h. If no coagulation in any degree is observed, the sample meets the requirements of the test for the absence of *Staphylococcus aureus*.

B-8.6 Validity of the Tests for Total Aerobic Microbial Count

Grow the following test strains separately in tubes containing fluid soyabean casein digest medium at 30 °C to 35 °C for 18 h to 24 h or, for *Candida albicans*, at 20 °C for 48 h.

Staphylococcus aureus (ATCC 6538; NCTC 10788)

Bacillus subtilis (ATCC 6633; NCIB 8054)

Escherichia coli (ATCC 8739; NCIB 8545)

Candida albicans (ATCC 2091; ATCC 10231)

Dilute portions of each of the cultures using buffered sodium chloride peptone solution pH 7.0 to make test suspensions containing about 100

viable microorganisms per ml. Use the suspension of each of the microorganisms separately as a control of the counting methods, in the presence and absence of the preparation being examined, if necessary.

A count for any of the test organisms differing by not more than a factor of 10 from the calculated value for the inoculum should be obtained. To test the sterility of the medium and of the diluent and the aseptic performance of the test, carry out the total aerobic microbial count method using sterile buffered sodium chloride peptone solution pH 7.0 as the test preparation. There should be no growth of microorganisms.

B-8.7 Validity of the Tests for Specified Microorganisms

Grow separately the test strains of *Staphylococcus aureus* and *Pseudomonas aeruginosa* in fluid soyabean casein digest medium and *Escherichia coli* and *Salmonella typhimurium* at 30 °C to 35 °C for 18 h to 24 h. Dilute portions of each of the

IS 19090 : 2024

cultures using buffered sodium chloride peptone solution pH 7.0 to make test suspensions containing about 10^3 viable microorganisms per ml. Mix equal volume of each suspension and use 0.4 ml (approximately 10^2 microorganisms of each strain) as an inoculum in the test for *E. coli*, *S.*

typhimurium, *P. aeruginosa* and *S. aureus*, in the presence and absence of the extract preparation being examined, if necessary. A positive result for the respective strain of microorganism should be obtained.

ANNEX C

(Foreword)

COMMITTEE COMPOSITION

Siddha Sectional Committee, AYD 05

<i>Organization</i>	<i>Representative(s)</i>
National Institute of Siddha, Chennai	PROF DR R. MEENAKUMARI (<i>Chairperson</i>)
Central Council for Research in Siddha, Chennai	DR SHYAMALA RAJKUMAR DR K SAMRAJ (<i>Alternate I</i>) DR V. AARTHI (<i>Alternate II</i>)
Central Drugs Standard Control Organisation, New Delhi	SHRI SUSHANT SHARMA DR RACHNA PALIWAL (<i>Alternate</i>)
CSIR-Central Institute of Medicinal and Aromatic Plants (CSIR-CIMAP), Lucknow	DR KARUNA SHANKER DR DEBABRATA CHANDA (<i>Alternate I</i>) DR KAPIL DEV (<i>Alternate II</i>)
CSIR-Traditional Knowledge Digital Library Unit (CSIR-TKDL), New Delhi	DR VIJAYALAKSHMI ASTHANA
Department of Pharmacy, Siddha Central Research Institute (CCRS-SCRI), Chennai	DR MS SHREE DEVI DR S.VINAYAK (<i>Alternate</i>)
Drug Testing Laboratory for Testing of A & S, (CCRAS – CSMCARI), Chennai	DR R. ILAVARASAN DR S. CHITRA (<i>Alternate I</i>) Ms MONIKA. N (<i>Alternate II</i>)
Government Siddha Medical College, Chennai	DR D. SASIKUMAR DR K. NALINA SARASWATHI (<i>Alternate</i>)
Government Siddha Medical College, Palayamkottai	DR T. KOMALAVALLI ALIAS MEHALA DR A. BALAMURUGAN (<i>Alternate I</i>) DR S. SIVANANDAN (<i>Alternate II</i>)
Kokila Siddha Hospital and Research Centre, Madurai	DR J. JEYAVENKATESH DR P. PAVITHRA (<i>Alternate</i>)
National Commission for Indian System of Medicine, New Delhi	DR M.V. MAHADEVAN DR N. J. MUTHUKUMAR (<i>Alternate I</i>) DR V. JASMINE RANJANA (<i>Alternate II</i>)
National Institute of Siddha, Chennai	DR S.SUDHA REVATHY DR V. SUBA (<i>Alternate I</i>) DR B. ANBARASAN (<i>Alternate II</i>)
National Medicinal Plants Board, New Delhi	DR R. MURUGESWARAN
Pharmacopoeia Commission for Indian Medicine & Homoeopathy, (PCIM&H), Ghaziabad	DR V. VIJAYAKUMAR DR NIKHIL M. JIRANKALGIKAR (<i>Alternate</i>)
State Drug Licensing Authority (Indian Medicine), Chennai	DR Y.R. MANEKSHAH DR G. SUBASH CHANDRAN (<i>Alternate</i>)
Tamil Nadu Ayurveda, Siddha Unani Manufacturers Association (TASUDMA), Chennai	DR L. SIVAKUMAR DR S. ILANGO (<i>Alternate I</i>) DR J. DINESH KUMAR (<i>Alternate II</i>)
Tamil Nadu Medicinal Plant Farms and Herbal Medicine Corporation Limited (TAMPCOL), Chennai	DR A. JAHEER HUSSAIN

IS 19090 : 2024

<i>Organization</i>	<i>Representative(s)</i>
The Indian Medical Practitioners' Co-operative Pharmacy & Stores (IMPCOPS), Chennai	DR R. KANNAN DR K. KADER MOHIDEEN (<i>Alternate I</i>) DR S. SUDARSHAN (<i>Alternate II</i>)
The Tamil Nadu Dr MGR Medical University, Chennai	DR N. KABILAN DR E. M. MANIKGANTAN (<i>Alternate</i>)
BIS Directorate General	SHRI UNNIKRIISHNAN A. R., SCIENTIST 'G' AND HEAD (AYUSH) [REPRESENTING DIRECTOR GENERAL (<i>Ex-officio</i>)]

Member Secretary
DR KIRUTHIGA G
SCIENTIST 'B'/ASSISTANT DIRECTOR
(AYUSH), BIS

Panel for Raw drugs Medicinal Plants, AYD 05/Panel 04

<i>Organization</i>	<i>Representative(s)</i>
National Institute of Siddha, Chennai	DR SUDHA REVATHY (<i>Convener</i>)
Central Council for Research in Ayurvedic Sciences, New Delhi	DR RAGHAVENDRA NAIK
Pharmacopoeia Commission for Indian Medicine & Homoeopathy, Ghaziabad	DR A. JAYANTHY
Central Council for Research in Unani Medicine, New Delhi	DR R. P. MEENA

Bureau of Indian Standards

BIS is a statutory institution established under the *Bureau of Indian Standards Act, 2016* to promote harmonious development of the activities of standardization, marking and quality certification of goods and attending to connected matters in the country.

Copyright

BIS has the copyright of all its publications. No part of these publications may be reproduced in any form without the prior permission in writing of BIS. This does not preclude the free use, in the course of implementing the standard, of necessary details, such as symbols and sizes, type or grade designations. Enquiries relating to copyright be addressed to the Head (Publication & Sales), BIS.

Review of Indian Standards

Amendments are issued to standards as the need arises on the basis of comments. Standards are also reviewed periodically; a standard along with amendments is reaffirmed when such review indicates that no changes are needed; if the review indicates that changes are needed, it is taken up for revision. Users of Indian Standards should ascertain that they are in possession of the latest amendments or edition by referring to the website-www.bis.gov.in or www.standardsbis.in.

This Indian Standard has been developed from Doc No.: AYD 05 (24706).

Amendments Issued Since Publication

Amend No.	Date of Issue	Text Affected

BUREAU OF INDIAN STANDARDS

Headquarters:

Manak Bhavan, 9 Bahadur Shah Zafar Marg, New Delhi 110002

Telephones: 2323 0131, 2323 3375, 2323 9402

Website: www.bis.gov.in

Regional Offices:

	Telephones
Central : 601/A, Konnectus Tower -1, 6 th Floor, DMRC Building, Bhavbhuti Marg, New Delhi 110002	{ 2323 7617
Eastern : 8 th Floor, Plot No 7/7 & 7/8, CP Block, Sector V, Salt Lake, Kolkata, West Bengal 700091	{ 2367 0012 2320 9474
Northern : Plot No. 4-A, Sector 27-B, Madhya Marg, Chandigarh 160019	{ 265 9930
Southern : C.I.T. Campus, IV Cross Road, Taramani, Chennai 600113	{ 2254 1442 2254 1216
Western : 5 th Floor/MTNL CETTM, Technology Street, Hiranandani Gardens, Powai Mumbai 400076	{ 25700030 25702715

Branches : AHMEDABAD, BENGALURU, BHOPAL, BHUBANESHWAR, CHANDIGARH, CHENNAI, COIMBATORE, DEHRADUN, DELHI, FARIDABAD, GHAZIABAD, GUWAHATI, HARYANA (CHANDIGARH), HUBLI, HYDERABAD, JAIPUR, JAMMU, JAMSHEDPUR, KOCHI, KOLKATA, LUCKNOW, MADURAI, MUMBAI, NAGPUR, NOIDA, PARWANOO, PATNA, PUNE, RAIPUR, RAJKOT, SURAT, VIJAYAWADA.