BUREAU OF INDIAN STANDARDS

BANGALORE BRANCH LABORATORY

13/06/2016

Subject: Comments on IS 4276:2014

This bears reference to the Indian Standard Specifications for edible oils published in 2014. While going through IS 4276:2014, following observations have been made.

S.	Test requirement / Ref. cl.	Comments
No. 1.	5.4 Oils shall not contain aflatoxin, more than 30 µg/kg, when tested by the method prescribed in IS/ISO 14718 or as prescribed in Annex A.	 Although the requirement for aflatoxin is given as quantitative (30 μg/kg), the method given in Annex A of relevant ISS appears to be for checking the presence or absence. Interpretation of results (Negative or positive) as given in cl.A-3.1 is not clear.
2	5.8.1.2 The product shall not contain aflatoxin, more than 5 μg/kg, when tested by the method prescribed in IS/ISO 14718 or as prescribed in Annex A.	do
3.	5.7.Table 1(ix), Hexane, in case of Refined Soybean Oil	 The requirement for Hexane in the ISSs given is 5. 00 ppm (mg/kg), Max. But as per last sentence of cl.B-1 (Annex B), the test method is suitable for determination of quantities of hexane between 10 and 1 500 mg/kg in fats and oils. It is proposed the source of availability of reference vegetable oil, solvent free, cl.B-3.4 (required for spiking with hexane for determination of calibration factor) may also be specified in the ISSs. (Pl. see Note). As per cl.B-6.1 Determination of the Calibration Factor, in the table (μl/5g v/s mg/100g), for the volume (μl) of Hexane added per 5 g of reference oil, the respective concentration (mg/100g) of Hexane in refined oil (spiked) does not appear to be correct. Values in given table are: Spiking of reference oil with Technical Hexane μl/5g 0.5 1 2 4 7 10 mg/100g 67 134 268 536 938 1340
		Spiking of reference oil with n-Hexane Mathematical Processing Content of Table 1988 1940

πg/100g 0.5 1 2 4 7 10 mg/100g 6.7 13.4 26.8 53.6 93.8 134.0
Spiking of reference oil with n-Hexane
μl/5g 0.5 1 2 4 7 10 mg/100g 6.6 13.2 26.4 52.8 92.4 132.0
It may please be noted that the values in 2ndrow when converted to mg/kg or ppm (i order to align the same in accordance with the requirements of edible oil standards) wi be as follow
Spiking of reference oil with Technical Hexane $\frac{\mu l/5g}{\mu l/5g} = 0.5 + 1 + 2 + 4 + 7 + 10$ $\frac{l/5g}{mg/lsg(nm)} = 124 + 268 + 526 + 028 + 1240$
mg/kg(ppm)67 134 268 536 938 1340 Spiking of reference oil with n-Hexane
μl/5g 0.5 1 2 4 7 10 mg/kg(PPM) 66 132 264 528 924 1320
The concentrations (mg/kg or ppm) prepared for making calibration curve armuch more than the specified requirements (5.00 ppm or 5.00 mg/kg).

Since, edible oils are likely to be covered under PCS of BIS, Competent Authority may kindly consider the above facts for making necessary changes/amendment in the specifications as mentioned above.

S. K. Arora Sc-B, OIC (QA & LRS)