BUREAU OF INDIAN STANDARDS

BANGALORE BRANCH LABORATORY

13/06/2016

Subject: Comments on IS 8361:2014

This bears reference to the Indian Standard Specifications for edible oils published in 2014. While going through IS 8361:2014, following observations have been made.

S.	Test requirement	Comments
No.	/ Ref. cl.	
1.	5.4 Oils shall not	1. Although the requirement for aflatoxin is given as quantitative (30 µg/kg), the method
	contain aflatoxin, more than 30	given in Annex A of relevant ISS appears to be for checking the presence or absence. 2. Interpretation of results (Negative or positive) as given in cl.A-3.1 is not clear.
		2. Interpretation of results (Negative or positive) as given in cl.A-3.1 is not clear.
	μg/kg, when tested by the	
	method	
	prescribed in	
	IS/ISO 14718 or	
	as prescribed in	
	Annex A.	
2	5.8.1.2 The	do
-	product shall not	
	contain aflatoxin,	
	more than 5	
	μg/kg, when	
	tested by the	
	method	
	prescribed in	
	IS/ISO 14718 or	
	as prescribed in	
	Annex A.	
3.	5.7. Table 1(x), Hexane, in case	1. The requirement for Hexane in the ISSs given is 5. 00 ppm (mg/kg), Max. But as per last sentence of cl.B-1 (Annex B), the test method is suitable for determination of
	of Refined	quantities of hexane between 10 and 1 500 mg/kg in fats and oils.
	Palmolein	2. It is proposed the source of availability of reference vegetable oil, solvent free, cl.B-
		3.4 (required for spiking with hexane for determination of calibration factor) may
		also be specified in the ISSs. (Pl. see Note).
		3. As per cl.B-6.1 Determination of the Calibration Factor, in the table (μ l/5g v/s
		mg/100g), for the volume (μl) of Hexane added per 5 g of reference oil, the respective
		concentration (mg/100g) of Hexane in refined oil (spiked) does not appear to be correct.
		Values in given table are:
		Spiking of reference oil with Technical Hexane
		Spaning of foreigned on what foreigned
		$\frac{1}{\mu l/5g}$ 0.5 1 2 4 7 10
		mg/100g 67 134 268 536 938 1340
		Spiking of reference oil with n-Hexane
		$\frac{1}{\mu l/5g}$ 0.5 1 2 4 7 10
		mg/100g 66 132 264 528 924 1320
		However, correct values in 2 nd row of table expressed as mg/100 g should be:
		Spiking of reference oil with Technical Hexane

πg/100g 0.5 1 2 4 7 10 mg/100g 6.7 13.4 26.8 53.6 93.8 134.0
Spiking of reference oil with n-Hexane
μl/5g 0.5 1 2 4 7 10 mg/100g 6.6 13.2 26.4 52.8 92.4 132.0
It may please be noted that the values in 2ndrow when converted to mg/kg or ppm (in order to align the same in accordance with the requirements of edible oil standards) will be as follow
Spiking of reference oil with Technical Hexane μ1/5g 0.5 1 2 4 7 10 mg/kg(ppm)67 134 268 536 938 1340
Spiking of reference oil with n-Hexane
μl/5g 0.5 1 2 4 7 10 mg/kg(PPM) 66 132 264 528 924 1320
The concentrations (mg/kg or ppm) prepared for making calibration curve are much more than the specified requirements (5.00 ppm or 5.00 mg/kg).

Since, edible oils are likely to be covered under PCS of BIS, Competent Authority may kindly consider the above facts for making necessary changes/amendment in the specifications as mentioned above.

S. K. Arora Sc-B, OIC (QA & LRS)